

## DEFENSE AGAINST THE SUGARCANE APHID www.sorghumcheckoff.com

# STAND AGAINST THE PEST

- SUGARCANE APHID ORIGIN & HOST PLANTS
  - IDENTIFYING THE PEST 5
  - BIOLOGY OF THE SUGARCANE APHID 6
    - APHID EFFECTS: HONEYDEW 7
    - APHID EFFECTS: SOOTY MOLD 8
    - APHID EFFECTS: LEAF NECROSIS
      - SCOUTING FOR THE PEST 10
      - WHEN TO TREAT FOR APHIDS 11
        - INSECTICIDE OPTIONS 12
        - INSECTICIDE APPLICATION 13
  - YIELD LOSS IF LEFT UNCONTROLLED 14
    - REASONS FOR POOR CONTROL 14
      - HYBRIDS AND RESISTANCE 15
        - RESISTANCE RESEARCH 15
      - HARVEST CONSIDERATIONS 16
        - ADDITIONAL RESOURCES 17
    - BEST MANAGEMENT PRACTICES 18

#### SUGARCANE APHID IN THE U.S.

The sugarcane aphid, *Melanaphis sacchari*, was discovered late in the 2013 growing season in the coastal sorghum regions. The sugarcane aphid is a new pest to the U.S. sorghum industry, and it is capable of causing substantial damage to the crop if left unmanaged. It is important for producers to be proactive and constantly scout for and monitor the pest because early detection is critical to minimize the aphid's harmful effects.

#### HOST PLANTS



Sugarcane aphids will only survive and multiply significantly in sorghum genotypes, including Johnsongrass, shattercane, sorghum sudangrass, sudangrass, forage sorghum and grain sorghum.

Sugarcane aphids will not survive on any of the major

crops such as corn, cotton, soybeans or wheat. However, sugarcane aphids are distributed by the wind, so small colonies can be found in these crops, but at this time, they have not been found to be of economic importance.

#### **IDENTIFYING THE PEST**



Identifying and treating for the sugarcane aphid in a timely manner is crucial. Sugarcane aphids are **gray**, **tan** or **pale yellow**, often called the white aphid. A pair of **black cornicles** or tailpipes can be seen on the aphid's back end. **Tips of feet** 

and antennae are black. A magnifying glass is required to see the black cornicles, feet and antennae. Colonies, or groups of sugarcane aphids are usually clustered around the midrib of the bottom side of the sorghum leaf. Additionally, sugarcane aphids are prolific producers of honeydew. The honeydew often originates from the aphids on the under side of leaves directly above.



#### BIOLOGY OF THE SUGARCANE APHID

1

All sugarcane aphids are females and are born pregnant. During their lifespan, aphids average approximately three live births a day.

2

Sugarcane aphid offspring become adults within five days and live an average of 28 days.

- 3 A proportion of the aphids in a colony will develop wings as adults. These winged sugarcane aphids are then dispersed with wind currents.
- 4

They have piercing, sucking mouth parts and feed on the sap of the plant, taking away nutrients that would otherwise go to growth and grain production.

5

Sugarcane aphids excrete a honeydew substance that causes the leaves to be shiny and sticky.

6

These pests are known to overwinter on sorghum species as far north as central Texas. It is currently unknown how far north the sugarcane aphid can potentially overwinter, but it is doubtful they can overwinter very far north.

#### **APHID EFFECTS: HONEYDEW**



Heavy infestations of the sugarcane aphid cause leaves to be covered with a **sticky**, **shiny** substance called **honeydew**. This honeydew is made up of plant sugars and water, which are harmless to animals. Honeydew is water soluble and washes off of the plant by rain or sprinkler irrigation. If honeydew is left on the plant, it eventually dries into a harmless residue.

#### APHID EFFECTS: SOOTY MOLD



**Black sooty mold** will often begin growing on the honeydew of sorghum leaves. This black sooty mold blocks out sunlight and decreases the plant's ability to produce sugars through photosynthesis. Over time, black sooty mold causes leaves to turn yellow and eventually the leaf tissue dies.

#### **APHID EFFECTS: LEAF NECROSIS**



Loss of plant sap from the sugarcane aphid feeding on sorghum leaves takes away nutrients from the plant that could otherwise be utilized for plant health and grain yield. Plant stress caused by the sugarcane aphids can also lead to uneven and lack of head emergence, poor grain set and will likely contribute to an increase in lodging.

#### SCOUTING FOR THE PEST



Fields should be scouted once a week for signs of the aphid. Once they are found, begin scouting at least twice a week.

P Honeydew found on the lower leaves in small hot spots within a field is often the first sign of sugarcane aphids.

**3** To determine if an insecticide treatment is needed, note the presence and number of aphids on leaves in at least four areas of the field.

In each area of the field examined, collect a leaf from the bottom and top of 20 plants and observe the presence and number of aphids.

#### WHEN TO TREAT FOR APHIDS

Threshold levels change and vary between regions. It is important to check with local experts to determine regional thresholds and scouting procedures. In general, two methods are currently being utilized to determine when to apply an insecticide to treat for sugarcane aphids.

Texas A&M University entomologists recommend threshold has been reached when 25 percent of the plants are infested with 50 aphids per leaf.

Mississippi State has explored thresholds by growth stage, which are described in the following table. **Regardless of the threshold level used, it is critical that insecticide application occurs as soon as possible.** Otherwise, severe loss can occur.

Growth Stage	Threshold
Pre-boot & Boot	20 percent of plants infested with localized area of heavy honeydew and established aphid colonies.
Heading, Soft Dough & Dough	30 percent of plants infested with localized area of heavy honeydew and established aphid colonies.
Black Layer	Heavy honeydew and established aphid colonies on the flag leaf or in the head. Only treat to prevent harvest problems. Observe preharvest insecticide intervals.

#### **INSECTICIDE OPTIONS**



As of October, 2021, there are three effective insecticide options available to control sugarcane aphid. These are:

**Sivanto Prime (flupyradifurone)**, sold by Bayer CropScience

Transform (sulfoxaflor), sold by Corteva Sefina (inscalis or afidopyropen), sold by BASF.

All three products are not harmful to beneficial insects, with are crucial in maintaining control of sugarcane aphids. **Care should be taken in adding other insecticides to the mix that could be harmful to beneficial insects**. Additionally, movement of the insecticides in the plant is minimal, making spray coverage essential for adequate control.

### **INSECTICIDE APPLICATION**

- When applying insecticides, it is important to read the full product label and follow instructions carefully.
- In general, ground rig application tends to work better than air application.
- Insecticide coverage is critical for sugarcane aphid control. Use spray nozzles and a spray pressure that maximizes coverage.
- Increasing the volume of water above the minimum requirements listed on the insecticide labels is recommended.
- Control will be lessened if insecticide application is made under cool conditions, including temperatures into the low 60 degrees Fahrenheit.
- The duration of control will vary from 7 to 28 days, depend on the insecticide and rate used, the initial control level achieved and the intensity of new infestations entering the field.
- Check with local experts on the use of adjuvants. Research on their effectiveness has been mixed.

#### YIELD LOSS IF LEFT UNCONTROLLED

Crop Stage at 20% Infestation	Percent Yield Loss with no Treatment
Pre-boot	81-100%
Boot	52-69%
Heading	67%
Soft Dough	21%

Source: Mississippi State University

#### REASONS FOR POOR CONTROL

Several factors can impact the effectiveness of insecticides in controlling the sugarcane aphid. Here are a few potential reasons for poor control:

- Waiting too long to make the insecticide application and the number of sugarcane aphids are simply overwhelming.
- Rain occurs soon after application, washing insecticide off of the leaf.
- Cool temperatures occurring at or soon after application causing sugarcane aphids to feed less and thus not ingest insecticide.
- Poor plant coverage, usually a result of not using enough water.
- Not using the appropriate insecticide rate.

#### **HYBRIDS AND TOLERANCE**

Hybrids with tolerance to sugarcane aphid are readily available from most seed companies. Tolerant hybrids are not immune to the sugarcane aphid, but the rate of aphid increase is much slower compared to susceptible hybrids.

In many cases tolerant hybrids will not need an insecticide application. If threshold level is reached then generally a single insecticide application is all that will be needed during the season. Under severe regional infestation levels susceptible hybrids often require multiple insecticide application.

#### **RESISTANCE RESEARCH**

Private companies, university and USDA-ARS breeders are working to discover new sources for improved resistance to sugarcane aphids. In addition, companies are constantly screening there new hybrids for their level of resistance/ tolerance. In choosing a hybrid, check the seed companies rating for sugarcane aphid tolerance. The level of tolerance does matter!

#### HARVEST CONSIDERATIONS



If sugarcane aphids are in significant numbers in the sorghum head or on the upper leaves, harvesting can be impeded due to honeydew. There is not a good threshold level at this time for when to spray sugarcane aphids to assist in harvest.

If sugarcane aphids are present and produce a significant amount of honeydew that can gum up the combine, many producers are opting to spray an insecticide to eliminate the risk of harvesting issues. Sivanto Prime, Transform and Sefina insecticides all have a preharvest interval of 14 days. If desired, these can be mixed with harvest-aid products such as glyphosate or sodium chlorate to facilitate plant dry down.

#### **REGIONAL EXPERTS**

\*This is not an inclusive list. University of Arkansas Extension | Gus Lorenz glorenz@uaex.edu Kansas State University | Brian McCornack mccornac@ksu.edu Louisiana State University AgCenter | Sebe Brown SBrown@agcenter.lsu.edu Mississippi State University | Angus Catchot acatchot@entomology.msstate.edu Oklahoma State University | Tom Royer tom.royer@okstate.edu Sorghum Checkoff | Brent Bean

brentb@sorghumcheckoff.com

Texas A&M AgriLife Extension | David Kerns dlkerns@ag.tamu.edu

#### **ADDITIONAL RESOURCES**

Sorghum Checkoff

www.sorghumcheckoff.com

#### Kansas State

Sugarcane Aphid- Scouting and Threshold

Texas A&M Agrilife Extension Managing Insect and Mite Pests of TX Sorghum

> Mississippi State Extension Sugarcane Aphid Management

#### BEST MANAGEMENT PRACTICES

Sugarcane aphids can only survive on sorghum-related species. To help reduce local populations, control Johnsongrass, volunteer sorghum and other sorghum species in and around your fields during winter and spring prior to planting.

2

Consider planting a hybrid that has shown tolerance to the aphid. Tolerance does not mean immunity to the aphid. These hybrids still require monitoring and treating with an insecticide if action thresholds are reached. Be careful in giving up hybrid adaptability, yield potential and other favorable agronomic characteristics. In most cases, these characteristics should not be given up in order to plant a sugarcane aphid tolerant hybrid.

3

Plant seeds treated with an insecticide seed treatment. These seed treatments will protect sorghum from potential early season infestations. Acceptable seed treatments include Cruiser (thiamethoxam), Poncho and Nipsit (clothianidin) and Gaucho (imidaclorprid).

4

**Plant early**. In many areas of the country, sugarcane aphids tend to infest fields later in the growing season. Early planting may avoid infestation.

# 5 Scout fields early and often using proper procedures to determine the level of aphid infestation. Once sugarcane aphid infestation occurs in the field, the number of aphids can increase guickly.

6

Apply insecticide as soon as the action threshold is reached. Threshold levels change and vary with individual states. Check with your local experts for current information. In general, the threshold is reached when 25 percent of the plants are infested with 50 aphids per leaf.

7

Use only recommended insecticides and follow label rates and application instructions. Coverage is critical. Best results are achieved when high volumes of water are used.

**Preharvest.** If aphids are present in the upper canopy or grain panicle in sufficient numbers to produce honeydew, consider applying an insecticide in order to prevent potential issues with harvest. If a harvestaid product is used, tank mixing with the insecticide has worked well when the sugarcane aphid is present.

Avoid use of insecticides, especially pyrethroids, that are harmful to beneficial insects because they may result in sugarcane aphid numbers increasing rapidly.

### TAKE CONTROL WIN PROFITABILITY SORGHUN HEART WWW.sorghumcheckoff.com