



U.S. GRAINS
COUNCIL

2024/2025
SORGHUM
QUALITY REPORT



As a USDA programs participant, the U.S. Grains Council is committed to complying with non-discrimination policies from federal, state and local civil rights laws and those of the USDA. Visit the USDA website page (<https://www.usda.gov/non-discrimination-statement>) for details.

Introductionii

Methodology.....1

Quality Test Results2

 A. Grade Factors2


 B. Tannins7

 C. Chemical Composition.....8

 D. Physical Factors10

U.S. Sorghum Supporting Information12

U.S. Grains Council Contact Information (back cover)



The U.S. Grains Council (Council) is pleased to present the findings from its *2024/2025 Sorghum Quality Report*. This is the sixth annual edition of a report designed to provide international customers and other interested parties with accurate, unbiased information about the quality of the U.S. sorghum crop. A total of 95 samples was collected from the outbound shipments of fifteen country elevators and analyzed for the grade factors established by the U.S. Department of Agriculture as well as chemical composition and other quality characteristics not reported elsewhere. The results are summarized at the U.S. Aggregate level.

The Council's mission is one of developing markets, enabling trade and improving lives. To help fulfill this mission, the Council is pleased to offer this report as a service to our partners. We hope it provides valuable information about the quality of U.S. sorghum to our valued trade partners.




Sorghum production in the United States can be grouped into the two main harvest areas shown in the map below (the Early Harvest Area and the Late Harvest Area). For this *2024/2025 Sorghum Quality Report*, the target population was sorghum from these two key U.S. sorghum-producing areas representing nearly 100% of the U.S. sorghum exports.¹ The results reflect the quality of blended samples pulled from the outbound sorghum shipments of U.S. elevators.

The targeted number of samples was proportionately stratified according to each sorghum-producing area's share of total U.S. sorghum exports, as shown on the map.

The Council recruited participating elevators across these two areas by email or phone. Participants were instructed to collect 2,200 grams (about five pounds) of sorghum per sample bag provided. Samples were requested to be collected from individual outbound rail or truck shipments as the sorghum was being loaded or from inventories expected to be shipped for export.

SORGHUM HARVEST AREAS



- A total of 95 blended sorghum samples from fifteen participating elevators was received from August 5, 2024, through February 18, 2025. Samples were received by the Amarillo Grain Exchange, Inc. (AGE) in Amarillo, Texas, and tested for the sorghum grade factors. Following grade factor testing, the samples were then sent to the Grain Quality Testing Lab at Kansas State University's Grain Science Department in Manhattan, Kansas, for chemical composition and physical factor analysis.
- Averages and standard deviations for each quality factor were calculated for the Early Harvest Area and the Late Harvest Area. The U.S. Aggregate average and standard deviation were then calculated based on each Harvest Area's proportion of total U.S. sorghum exports following standard statistical techniques for proportionate stratified sampling. The results are reported only for the U.S. Aggregate and not the individual Harvest Areas.

¹Source: USDA/NASS and Centrec estimates

A. GRADE FACTORS

The U.S. Department of Agriculture's Federal Grain Inspection Service (FGIS) has established numerical grades, definitions and standards for grains. The attributes that determine the numerical grades for sorghum are test weight, broken kernels and foreign material (BNFM), foreign material, total damage and heat damage.

Two of the 95 total samples collected for this year's report weighed less than 1,000 grams. This prohibited test weight analysis and resulted in 93 samples analyzed for test weight.

SUMMARY: GRADE FACTORS

2024/2025						2023/2024			2022/2023			5-Year Average 19/20-23/24	
	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	No. of Samples	Avg.	Std. Dev.	Avg.	Std. Dev.
U.S. Aggregate						U.S. Aggregate			U.S. Aggregate			U.S. Aggregate	
Test Weight (lb/bu)	93	57.7	2.26	52.1	62.3	104	58.4*	1.75	96	58.8*	2.03	58.7*	1.77
Test Weight (kg/hl)	93	72.0	2.82	65.0	77.7	104	75.2*	2.25	96	75.7*	2.61	75.6*	2.27
BNFM (%)	95	1.9	1.11	0.1	5.4	104	1.8	0.93	96	1.4*	1.15	1.6*	0.95
Foreign Material (%)	95	0.8	0.60	0.0	3.6	104	0.9	0.66	97	0.7	0.67	0.7*	0.54
Total Damage (%)	95	0.3	0.44	0.0	4.3	104	0.0*	0.00	97	0.0	0.00	0.1*	0.26
Heat Damage (%)	95	0.0	0.00	0.0	0.0	104	0.0	0.00	97	0.0	0.00	0.0	0.00

*Indicates average was significantly different from current year's sorghum, based on a 2-tailed t-test at the 95.0% level of significance.


TEST WEIGHT

Test weight is defined as the weight of grain required to fill a specific volume (Winchester bushel). This measure of bulk density is often used as a general indicator of overall quality and as a gauge of endosperm hardness for value-added processing.

The test involves filling a test cup of known volume through a funnel held at a specific height above the test cup to the point where grain begins to pour over the sides of the test cup. A strike-off stick is used to level the grain in the test cup, and the grain remaining in the cup is weighed. The weight is then converted to and reported in the traditional U.S. unit, pounds per bushel (lb/bu). Results are also reported in kilograms per hectoliter (kg/hl) in this report. The average and standard deviation for the 2024/2025 samples were the following:

- Average: **57.7 lb/bu (72.0 kg/hl)**, lower than 2023/2024 (58.4 lb/bu or 75.2 kg/hl) and the 5YA¹ (58.7 lb/bu or 75.6 kg/hl) but higher than the minimum for U.S. No. 1 grade (57.0 lb/bu or 73.4 kg/hl)
- Standard deviation: **2.26 lb/bu (2.82 kg/hl)**, higher than 2023/2024 (1.75 lb/bu or 2.25 kg/hl) and the 5YA (1.77 lb/bu or 2.27 kg/hl)

TEST WEIGHT (lb/bu & kg/hl)
U.S. AGGREGATE RESULTS SUMMARY



¹The 5YA represents the simple average of the quality factors' average or standard deviation from the 2019/2020, 2020/2021, 2021/2022, 2022/2023 and 2023/2024 Sorghum Quality Reports.


BROKEN KERNELS AND FOREIGN MATERIAL

Broken kernels and foreign material (BNFM), an indicator of the amount of clean, sound sorghum available for feed and processing, is reported as the sum of broken kernels as a percent of the dockage-free sample weight and the foreign material.

Broken kernels is defined as all material that passes through a 5/64th-inch triangular-hole sieve and over a 2.5/64th-inch round-hole sieve.

Foreign material is defined as all material, except sorghum, that remains on top of the 5/64th-inch triangular-hole sieve and all matter other than sorghum, which passes over a No. 6 riddle. Foreign material is reported as a sum of the mechanically-separated foreign material as a percent of the dockage-free sample weight and the handpicked foreign material as a percent of the handpicked sample portion weight. The average and standard deviation for the 2024/2025 samples were the following:


- Average: **1.9%**, similar to 2023/2024 (1.8%), higher than the 5YA (1.6%) but lower than the maximum for U.S. No. 1 grade (3.0%)
- Standard deviation: **1.11%**, higher than 2023/2024 (0.93%) and the 5YA (0.95%)



FOREIGN MATERIAL

Foreign material, a part of BNFM, consists of non-sorghum material and dust. Foreign material is generally higher in moisture content than the sorghum itself and reduces feed and processing value. The average and standard deviation for the 2024/2025 samples were the following:


- Average: **0.8%**, similar to 2023/2024 (0.9%), higher than the 5YA (0.7%) but below the maximum for U.S. No. 1 grade (1.0%)
- Standard deviation: **0.60%**, similar to 2023/2024 (0.66%) and the 5YA (0.54%)



TOTAL DAMAGE

Total damage is the percentage of kernels and pieces of kernels that are damaged in some way. A representative working sample of 15 grams of BNFM-free sorghum is visually examined by a properly trained individual for the content of damaged kernels. Types of damage include germ-damaged kernels, ground- or weather-damaged kernels, diseased kernels, frost-damaged kernels, heat-damaged kernels, insect-bored kernels, mold-damaged kernels (surface or internal), mold-like substance, purple-pigment-damaged kernels and sprout-damaged kernels. Total damage is reported as the weight percentage of the working sample that is total damaged grain.

- Average: **0.3%**, higher than 2023/2024 (0.0%) and the 5YA (0.1%) but below the maximum for U.S. No. 1 grade (2.0%)
- Standard deviation: **0.44%**, higher than 2023/2024 (0.00%) and the 5YA (0.26%)



HEAT DAMAGE

Heat damage is a subset of total damage and consists of kernels and pieces of sorghum kernels that are materially discolored and damaged by heat. It occurs during heated air drying or in storage. Heat-damaged kernels are determined by a properly trained individual visually inspecting a 15-gram sample of BNFM-free sorghum. Heat damage, if found, is reported separately from total damage. The average for the 2024/2025 samples was the following:

- Average: **0.0%**, no heat damage was observed, the same as in the previous five years


B. TANNINS

Tannins are present in sorghum varieties that have a pigmented testa within their kernels. Tannins affect nutritional and functional properties as a result of interactions of the tannins in sorghum-containing rations. Values near or below 4.0 mg catechin equivalents (CE) per gram (g) sample by this method generally imply the absence of condensed tannins. Type III tannin sorghums usually have values greater than 8.0 mg CE/g.

Leucoanthocyanidins (catechins) and proanthocyanidins (tannins) are a class of flavonoids known as flavanols that react with vanillin in the presence of mineral acids to produce a red color. Vanillin reacts with flavanols, but other flavonoid compounds can give specific color development. The test involves grinding approximately 50 g of sound seed using a UDY grinder with a 1 mm sieve and accurately weighing 0.30 g of this sample for analysis. Extraction and analysis are performed using the vanillin-HCl test with blank subtraction to remove interference by sorghum pigments. Developed color is measured using a UV-Vis spectrophotometer at 500 nanometers. A standard curve is run using pure catechin. Tests are run in triplicates, and the average value is reported as mg CE/g sample on a dry basis.

Four of the 95 total samples collected and analyzed for grade factors at AGE were not received by Kansas State University, resulting in 91 samples analyzed for tannins.

- Tannin levels in all 91 samples tested were less than 4.0 mg CE/g, implying an absence of tannins, the same as in 2023/2024 and 2022/2023.



C. CHEMICAL COMPOSITION

Chemical composition of sorghum is important because the components of protein, starch and oil are of significant interest to end-users. These attributes provide additional information related to nutritional value for livestock and poultry feeding and other processing uses of sorghum. Chemical composition tests for protein, oil and starch were conducted using an approximately 50-gram sample in a Perten DA 7250 Near-Infrared Reflectance (NIR) instrument. Results are reported on a dry basis (percent of non-water material).

Four of the 95 total samples collected and analyzed for grade factors at AGE were not received by Kansas State University, resulting in 91 samples analyzed for chemical composition.

SUMMARY: CHEMICAL FACTORS


	2024/2025					2023/2024			2022/2023			5-Year Average 19/20-23/24	
	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	No. of Samples	Avg.	Std. Dev.	Avg.	Std. Dev.
U.S. Aggregate						U.S. Aggregate			U.S. Aggregate			U.S. Aggregate	
Protein (Dry Basis %)	91	10.6	1.63	7.4	13.7	104	11.1*	0.79	97	11.0*	0.64	11.0*	0.77
Starch (Dry Basis %)	91	73.9	1.21	70.5	76.1	104	73.1*	0.88	97	72.2*	1.11	72.7*	1.04
Oil (Dry Basis %)	91	5.1	0.27	4.0	5.5	104	4.8*	0.16	97	4.7*	0.19	4.7*	0.23

*Indicates average was significantly different from current year's sorghum, based on a 2-tailed t-test at the 95.0% level of significance.

PROTEIN

Protein is very important for poultry and livestock feeding, as it supplies essential sulfur-containing amino acids and improves the feed conversion efficiency. Protein is usually inversely related to starch concentration. Results are reported on a dry basis. The average and standard deviation for the 2024/2025 samples were the following:

- Average: **10.6%**, lower than 2023/2024 (11.1%) and the 5YA (11.0%)
- Standard deviation: **1.63%**, higher than 2023/2024 (0.79%) and the 5YA (0.77%)






STARCH

Starch is related to metabolizable energy for livestock and poultry. High starch concentration is often indicative of good kernel maturation/filling conditions and reasonably moderate kernel densities. Results are reported on a dry basis. The average and standard deviation for the 2024/2025 samples were the following:


- Average: **73.9%**, higher than 2023/2024 (73.1%) and the 5YA (72.7%)
- Standard deviation: **1.21%**, higher than 2023/2024 (0.88%) and the 5YA (1.04%)



OIL

Oil is an essential component of poultry and livestock rations. It serves as an energy source, enables fat-soluble vitamins to be utilized and provides certain essential fatty acids. Oil may also be an important co-product of sorghum value-added processing. Results are reported on a dry basis. The average and standard deviation for the 2024/2025 samples were the following:

- Average: **5.1%**, higher than 2023/2024 (4.8%) and the 5YA (4.7%)
- Standard deviation: **0.27%**, higher than 2023/2024 (0.16%) but similar to the 5YA (0.23%)



D. PHYSICAL FACTORS

Physical factors include other quality attributes that are neither grading factors nor chemical composition. Tests for physical factors provide additional information about the processing characteristics of sorghum for various uses, as well as its storability and potential for breakage in handling.

Four of the 95 total samples collected and analyzed for grade factors at AGE were not received by Kansas State University, resulting in 91 samples analyzed for physical factors.

SUMMARY: PHYSICAL FACTORS

2024/2025						2023/2024			2022/2023			5-Year Average 19/20-23/24	
	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	No. of Samples	Avg.	Std. Dev.	Avg.	Std. Dev.
U.S. Aggregate						U.S. Aggregate			U.S. Aggregate			U.S. Aggregate	
Kernel Diameter (mm)	91	2.52	0.17	2.12	2.89	73	2.49	0.19	97	2.50	0.14	2.51	0.14
1000-Kernel Weight (g)	91	25.17	2.96	19.95	34.56	73	23.52*	3.79	97	24.76	3.14	25.02	2.97


*Indicates average was significantly different from current year's sorghum, based on a 2-tailed t-test at the 95.0% level of significance.

KERNEL DIAMETER

Kernel diameter directly correlates with kernel volume, affects material handling practices and may indicate the maturity of kernels.

Kernel diameter is measured using a Perten Single Kernel Characterization System (SKCS 4100). The instrument records the individual diameter of 300 seeds and calculates the average seed diameter in millimeters (mm). The average and standard deviation for the 2024/2025 samples were the following:

- Average: **2.52 mm**, similar to 2023/2024 (2.49 mm) and the 5YA (2.51 mm)
- Standard deviation: **0.17 mm**, similar to 2023/2024 (0.19 mm) and the 5YA (0.14 mm)




1000-KERNEL WEIGHT

1000-kernel weight (commonly referred to as TKW) is the weight for a fixed number of kernels and is reported in grams. Kernel volume (or size) can be inferred from TKW. As TKW increases or decreases, kernel volume will proportionally increase or decrease.

The TKW is determined from the average weight of 300 individual kernel replicates using the SKCS 4100. The instrument weighs each seed to the nearest 0.01 mg and automatically calculates the TKW based on the average weight of the 300 individual seeds. The average TKW is reported in grams. The average and standard deviation for the 2024/2025 samples were the following:

- Average: **25.17 g**, higher than 2023/2024 (23.52 g) but similar to the 5YA (25.02 g)
- Standard deviation: **2.96 g**, lower than 2023/2024 (3.79 g) but similar to the 5YA (2.97 g)



U.S. SORGHUM GRADES AND GRADE REQUIREMENTS

Grade	Minimum Test Weight per Bushel (Pounds)	Maximum Limits of			
		Damaged Kernels		Broken Kernels and Foreign Material	
		Heat Damaged (Percent)	Total (Percent)	Foreign Material (part of total) (Percent)	Total (Percent)
U.S. No. 1	57.0	0.2	2.0	1.0	3.0
U.S. No. 2	55.0	0.5	5.0	2.0	6.0
U.S. No. 3 ¹	53.0	1.0	10.0	3.0	8.0
U.S. No. 4	51.0	3.0	15.0	4.0	10.0

U.S. Sample Grade is sorghum that:

- (a) Does not meet the requirements for the grades U.S. Nos. 1, 2, 3, or 4; or
- (b) Contains 8 or more stones which have an aggregate weight in excess of 0.2 percent of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria spp.*), 2 or more castor beans (*Ricinus communis L.*), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 8 or more cockleburrs (*Xanthium spp.*) or similar seeds singly or in combination, 10 or more rodent pellets, bird droppings, or an equivalent quantity of other animal filth in 1,000 grams of sorghum, 11 or more pieces of other material from any combination of animal filth, castor beans, crotalaria seeds, glass, stones, unknown foreign substances, and cockleburrs; or
- (c) Has a musty, sour, or commercially objectionable foreign odor (except smut odor); or
- (d) Is badly weathered, heating or otherwise of distinctly low quality.

¹Sorghum which is distinctly discolored shall not grade any higher than U.S. No. 3.

Source: Code of Federal Regulations, Title 7, Part 810, Subpart D, United States Standards for Sorghum

U.S. AND METRIC CONVERSIONS

Sorghum Equivalents	Metric Equivalents
1 bushel = 56 pounds (25.40 kilograms)	1 pound = 0.4536 kg
39.368 bushels = 1 metric ton	1 hundredweight = 100 pounds or 45.36 kg
15.93 bushels/acre = 1 metric ton/hectare	1 metric ton = 2204.6 lbs
1 bushel/acre = 62.77 kilograms/hectare	1 metric ton = 1000 kg
1 bushel/acre = 0.6277 quintals/hectare	1 metric ton = 10 quintals
56 lbs/bushel = 72.08 kg/hectoliter	1 quintal = 100 kg
	1 hectare = 2.47 acres



U.S. GRAINS COUNCIL

A global network of professionals building worldwide demand
and developing markets for U.S. grains and ethanol.



HEADQUARTERS

20 F Street NW, Suite 900 • Washington, D.C., 20001

Phone: 202-789-0789 • Fax: 202-898-0522

Email: grains@grains.org • Website: grains.org

PEOPLE'S REPUBLIC OF CHINA: Beijing

Tel1: 011-86-10-6505-1314 • Tel2: 011-86-10-6505-2320
Fax: 011-86-10-6505-0236 • china@grains.org

JAPAN: Tokyo

Tel: 011-81-3-6206-1041 • Fax: 011-81-3-6205-4960
japan@grains.org • www.grainsjp.org

KOREA: Seoul

Tel: 011-82-2-720-1891 • Fax: 011-82-2-720-9008
seoul@grains.org

TAIWAN: Taipei

Tel: 011-886-2-2523-8801 • Fax: 011-886-2-2523-0149
taipei@grains.org

MIDDLE EAST, AFRICA AND EUROPE: Tunis

Tel: 011-216-71-191-640 • Fax: 011-216-71-191-650
tunis@grains.org

INDIA: New Delhi

Tel: +91-11-4603-6437 • usgcindia@grains.org

SOUTH EAST ASIA: Kuala Lumpur

Tel: 011-603-2789-3288 • sea-oceania@grains.org

LATIN AMERICA: Panama City

Tel: 011-507-315-1008 • lta@grains.org

MEXICO: Mexico City

Tel1: 011-52-55-5282-0244 • Tel2: 011-52-55-5282-0973
Tel3: 011-52-55-5282-0977 • mexico@grains.org